

The Magellanic Bridge: evidence for a population of X-ray binaries

V. A. McBride,¹★ A. J. Bird,¹ M. J. Coe,¹ L. J. Townsend,¹ R. H. D. Corbet^{2,3}
and F. Haberl⁴

¹*School of Physics & Astronomy, University of Southampton, Southampton SO17 1BJ*

²*University of Maryland, Baltimore County, MD 21250, USA*

³*CRESST/Mail Code 662, NASA Goddard Space Flight Center, Greenbelt, MD 20771, USA*

⁴*Max-Planck Institut für Extraterrestrische Physik, Giessenbachstraße, 85748 Garching, Germany*

Accepted 2009 December 7. Received 2009 November 4; in original form 2009 September 24

ABSTRACT

INTEGRAL observations of the Small Magellanic Cloud region have resulted in the serendipitous detection of two transient hard X-ray sources in the Magellanic Bridge. In this paper, we present the timing and spectral characteristics of these sources across the 2–100 keV energy range, which, in conjunction with their optical counterparts, demonstrate that they are high-mass X-ray binaries (HMXBs) in the Magellanic Bridge. Together with one previously known HMXB system, and three candidates, these sources represent an emerging population of X-ray binaries in the Bridge, probably initiated by tidally induced star formation as a result of the gravitational interaction between the Large and Small Magellanic Clouds.

Key words: binaries: general – Magellanic Clouds.

1 INTRODUCTION

The Magellanic Clouds present a nearby, easily observable laboratory in which to study minor galaxy interactions. In this system, the Magellanic Bridge is thought to be a product of the tidal interaction between the Large and Small Magellanic Clouds (LMC and SMC). It contains both gas and stellar components, with the stellar population first discovered by Irwin, Kunkel & Demers (1985) and comprising only a young (<300 Myr) stellar component (Harris 2007), which is thought to have formed in situ. One identified massive X-ray binary has been detected in the western bridge (Kahabka & Hilker 2005). Given the tidal disruption in this region, a number of star formation tracers are expected in this active region. The focus of this work is to identify high-mass X-ray binaries (HMXBs) as tracers of star formation in the Magellanic Bridge.

Multiwavelength studies of the SMC have shown that it contains a large number of X-ray binary pulsars – all but one of them in Be/X-ray binary systems (Coe et al. 2005, 2009). This preponderance of young systems is most likely explained by tidally triggered star formation precipitated by the most recent close approach of the SMC and LMC (Gardiner & Noguchi 1996). However, the most recent close approach of the SMC and LMC was ~200 Myr ago – much longer than the evolutionary time-scale of Be/X-ray binaries, which implies that either there has been a significant delay between the encounter of the SMC and LMC and the onset of star formation, or that subsequent waves of star formation have given rise to these Be/X-ray binaries (Harris & Zaritsky 2004).

These objects, being tracers of star formation (Grimm, Gilfanov & Sunyaev 2003), give direct insights into the star formation history of their host galaxies.

The LMC, by comparison, has a very different population at high energies: whereas the SMC population comprises almost exclusively Be/X-ray binaries, the population of the LMC has representatives from all members of the X-ray binary classes, including black hole systems, low-mass X-ray binaries, and Be and supergiant high-mass systems. With the mass of the LMC being ~10× that of the SMC, tidal interactions between the galaxies would have a much greater effect on the SMC, and this may be reflected by the differing stellar populations.

The high-energy population of the Magellanic Bridge is not well known, but extrapolation from optical wavelengths indicates that there should be many young stellar systems towards the western bridge (Harris 2007), resembling the population in the SMC. The work presented here is the first systematic study of this region at hard X-ray energies. The rest of the paper is structured as follows: Section 2 presents the observations and data analysis. In Section 3, the new transient objects and their characteristics are presented, while other candidate high-energy sources are discussed in Section 4. The discussion and conclusions are presented in Sections 5 and 6, respectively.

2 OBSERVATIONS AND ANALYSIS

The IBIS telescope (Ubertini et al. 2003) aboard *INTEGRAL*, which is optimized for an energy range of 15–200 keV and has a field of view of 30° × 30°, is uniquely suited to observing large sky areas for point sources. As part of a key programme monitoring campaign on

★E-mail: vanessa@soton.ac.uk

Table 1. *INTEGRAL* observation log.

Rev. number	MJD _{start}	Exposure (ks)
745	54788.7	37.7
746	54791.8	40.2
747	54794.8	37.0
748	54797.7	41.1
749	54800.8	39.7
750	54803.7	46.0
751	54806.7	49.2
752	54809.7	51.9
753	54812.7	50.0
754	54815.7	39.2
755	54818.7	50.6
756	54821.7	49.5
796	54941.4	69.8
797	54944.4	55.6
812	54989.2	49.5
813	54992.2	40.2
814	54995.2	49.3
815	54998.2	48.7
816	55001.2	52.9
817	55004.2	55.4
818	55007.2	31.5

the SMC and 47 Tuc, *INTEGRAL* observed the SMC and Magellanic Bridge for approximately 90 ks per satellite revolution (~ 3 days) from 2008 November 11 to 2009 June 25. The exposure on the Magellanic Bridge (Table 1) is, in general, much smaller than the 90 ks observation time, both due to the fact that the Bridge is a few degrees from the main pointing position, and that the figures for exposure time have been corrected for instrument dead time.

Individual pointings (science windows) were processed using the *INTEGRAL* Offline Science Analysis v.7.0 (osa, Goldwurm et al. 2003) and were mosaicked using the weighted mean of the flux in the 3–10 keV (JEM-X) and 15–35 keV (IBIS) energy ranges. Proprietary software was used to mosaic the observations from successive revolutions to improve the exposure and thereby the sensitivity to faint sources. Light curves in these energy bands were generated on science window (~ 2000 s) and revolution time-scales. The IBIS energy band was chosen to maximize the detection significance of

SMC X-1, and hence other SMC accreting X-ray pulsars, which have similar spectral shapes to SMC X-1 in this energy range. An IBIS mosaic of data from revolutions 753–756 in the 15–35 keV energy range is shown in Fig. 1.

3 NEW SOURCES

3.1 IGR J015712–7259

A new source, IGR J015712–7259 (Coe et al. 2008) was discovered in revolution 752. It showed transient behaviour with a maximum significance of 7.7σ from revolutions 812 to 814 in IBIS, corresponding to an average flux of $3.3 \pm 0.3 \times 10^{-11}$ erg cm $^{-2}$ s $^{-1}$ (1.4×10^{37} erg s $^{-1}$ at 60 kpc). The source was detected in JEM-X at an average flux of $1.6 \pm 0.5 \times 10^{-12}$ erg cm $^{-2}$ s $^{-1}$ in 3–10 keV. The IBIS light curve is shown in Fig. 2. Timely follow-up with *Swift*/XRT on 2008 December 20 (MJD 54820) allowed a precise determination of the source position: $1^{\text{h}}57^{\text{m}}16^{\text{s}}$, $-72^{\circ}58'32''.8$ (J2000.0) with a 90 per cent error circle of radius 3.8 arcsec. This identifies the source with a star (USNO-B1 0170-0064697; Monet et al. 2003) having *B* and *R* magnitudes of 15.48 and 15.51, respectively (see Fig. 3). A total of ~ 300 counts extracted from a region around the *Swift*/XRT source showed pulsations at ~ 11 s.

A 15 ks *RXTE* observation on 2008 December 24 revealed that the source was pulsating with a period of 11.57809 ± 0.00002 s (see Fig. 4). The hard X-ray behaviour is illustrated in the IBIS light curve (Fig. 2), the *RXTE* folded light curve (Fig. 5) and the broad-band spectrum of the source around MJD 54824 (Fig. 6). The combined *Swift*/XRT and IBIS spectrum can be adequately fit ($\chi^2_{\nu} = 1.05$ for 33 dof) with an absorbed, exponentially cut-off power law and a free constant factor between the two instruments to account for the fact that the soft and hard spectra were not observed simultaneously. The constant factor is ~ 0.3 , while the photon index is 0.4 ± 0.2 , and the folding energy is 8^{+5}_{-3} keV. The absorption has been fixed to the neutral density of hydrogen along the line of sight to the SMC: 6×10^{20} cm $^{-2}$. The photon index is similar to those determined for X-ray pulsars in the wing of the SMC (McGowan et al. 2007). The transient nature of the source, combined with the detected pulsations, hard spectrum and the optical magnitudes of the counterpart are strongly suggestive of a Be/X-ray binary.




Figure 1. IBIS 15–35 keV mosaic of the SMC and Magellanic Bridge from revolutions 753–756. The contours show the HI distribution in the SMC and Bridge, and are from Putman et al. (2003).




Figure 2. IBIS light curve of IGR J015712–7259 in the 15–35 keV energy band.




Figure 3. DSS II *R*-band image of a 10×10 arcmin² field around the source IGR J015712–7259, showing the IBIS 90 per cent error circle and with the inset showing an expanded view (indicated by the square) of the counterpart and the *Swift* XRT 90 per cent error circle. North is up and east is to the left.

3.2 SWIFT J0208.4–7428

Another source was discovered during revolution 756 (MJD 54821.7) in the IBIS map at a position of $02^{\text{h}}07^{\text{m}}09^{\text{s}}$, $-74^{\circ}28'07''$ with a 3.6 arcmin error circle. It reached a maximum significance of 7.1σ during revolutions 753–756, with a corresponding average flux of 2.5×10^{-11} erg cm⁻² s⁻¹ in the 15–35 keV band ($\sim 1 \times 10^{37}$ erg s⁻¹ at 60 kpc). The source was weakly detected with JEM-X, at a flux of $1.5 \pm 0.6 \times 10^{-12}$ erg cm⁻² s⁻¹. The light curve in the 15–35 keV band in Fig. 7 clearly shows the flux rising through the observation sequence.

Archival data from *Swift*/XRT showed that a source in the Magellanic Bridge was observed with *Swift* on MJDs 54764.5 and 54809.5. Analysis of the XRT data shows that no source is present within the field of view during the observation on MJD 54764.5, but during MJD 54809, a source is detected at




Figure 4. Periodogram of the *RXTE* observation of IGR J015712–7259 on MJD 54824.




Figure 5. *RXTE* 3–10 keV light curve of IGR J015712–7259 folded at the pulse period of 11.578 s.




Figure 6. Combined *Swift*/XRT and IBIS spectrum of IGR J015712–7259, showing the residuals as $\Delta\chi$ in the lower panel. The data are plotted as crosses, the model fit as the histogram, while the smooth curve shows the unfolded spectrum as plotted on the secondary *Y*-axis.

$02^{\text{h}}06^{\text{m}}45^{\text{s}}.7$, $-74^{\circ}27'46''.3$ (J2000.0) with a 90 per cent error circle of 4 arcsec. This source falls well within the 90 per cent error circle of the *INTEGRAL* source, and can be clearly distinguished from RX J0209.6–7427 (the only previously known HMXB in the Magellanic Bridge, see Fig. 8). These contemporaneous observations (see light curve in Fig. 7) provide strong evidence for another




Figure 7. IBIS light curve of SWIFT J0208.4–7428 (red line shows *Swift*/XRT detection).




Figure 8. DSS II *R*-band image of a 10×15 arcmin² field surrounding SWIFT J0208.4–7428, showing the 3.6 arcmin radius 90 per cent IBIS error circle, the nearby known HMXB, RX J0209.6–7427, and with the inset showing an expanded view of the square region along with the *Swift* XRT error circle. North is up and east is to the left.

new X-ray source in the Magellanic Bridge. Only one object falls within the *Swift*/XRT error circle and can be identified with a $B = 14.41$ and $V = 14.75$ star (Demers & Irwin 1991), for which the $B - V$ colour and magnitudes are consistent with an early-type dwarf at the distance of the SMC.

4 CANDIDATE SOURCES

IBIS 15–35 keV and JEM-X 3–10 keV maps on time-scales varying between single revolutions up to five consecutive revolutions were searched for excesses which could indicate potential new sources in the Magellanic Bridge. A list of candidates is supplied in Table 2. None of these candidate sources correlate with known *ROSAT* sources (Voges et al. 1999, 2000), but it is worth bearing in mind that the *ROSAT* area coverage in this part of the sky is not complete. With the large error circles on these objects, it is not yet possible to constrain the nature of these sources, and, although they may be background AGN, their transient nature makes it more likely that they are binary systems in the Magellanic Bridge.

5 DISCUSSION

Before this work, only a single massive X-ray binary system was known in the Magellanic Bridge (RX J0209.6–7427; Kahabka & Hilker 2005). So far, all X-ray sources discovered and identified in the Magellanic Bridge have been transient, with no persistent sources detected down to a luminosity of 2.5×10^{36} erg s^{−1} (15–35 keV). In this respect, the population of the Magellanic Bridge seems to resemble that of the SMC more closely than it does of the LMC. The Magellanic Bridge contains a significant population of young blue stars (Demers & Irwin 1991; Harris 2007). Harris (2007) showed that the distribution of blue stars is most dense towards the SMC side and that there appear to be very few old, red giant branch stars in the fields studied. As most HMXBs in the SMC are found in regions populated by young stars (Yokogawa et al. 2003), this seems to indicate that the trend will be to find more HMXBs on the SMC side of the Bridge, where the stellar population is younger. However, in our work, the exposure is by no means uniform across the Magellanic Bridge (see Fig. 9) and this introduces a selection effect whereby new HMXBs are selectively found where the exposure is highest, i.e. towards the SMC side of the Bridge. Thus, we cannot yet be confident that the population of X-ray binaries so far uncovered in the Bridge is indicative of the distribution in the Bridge.

The prevalence of a young stellar population, and lack of an older one, suggests that the stellar population of the Magellanic Bridge formed in situ, rather than being tidally extracted from the LMC and SMC. In addition to the HMXBs discovered here, the region also shows strong evidence for recent star formation with ~ 100 OB associations identified in the western Magellanic Bridge (Bica & Schmitt 1995) as well as molecular clouds (Mizuno et al. 2006), which act as fuel for ongoing star formation.

It is very unlikely that the HMXBs so far discovered in the Magellanic Bridge could have been ejected from the SMC through supernovae kicks. Not only would they require systemic velocities in excess of 200 km s^{−1} (where most SMC HMXBs have estimated systemic velocities of ~ 30 km s^{−1}, Coe 2005), but we would expect to find a distribution of HMXBs around the SMC rather than preferentially concentrated in the Bridge.

To estimate what fraction of a potential HMXB population in the Magellanic Bridge has been observed to date, we use an estimate of 1 Be + NS binary system per square degree, as obtained by Kahabka & Hilker (2005). This estimate is based on the observed stellar density of stars $> 2.5 M_{\odot}$ in the Magellanic Bridge (Gardiner & Hatzidimitriou 1992), corrected through the initial mass function to give the number of B stars per square degree. Through estimates of the Be/B fraction and an assumption of how many Be stars have a neutron star companion, these authors estimate the number of Be + NS systems as between 0.68 and 0.96 per square degree, dependent on the shape of the initial mass function. Given that the Magellanic Bridge spans an area of ~ 6 deg² on the sky, we may expect up to 6 Be + NS systems in the Magellanic Bridge, which is roughly consistent with the number of sources and potential candidates reported up until now, including this work. At this early stage, it is premature to attempt quantitative estimates of an upper limit to the potential number of HMXBs in the Magellanic Bridge. A prediction based on star formation rate (e.g. Grimm et al. 2003) is prone to significant uncertainties in both the X-ray luminosity function of this potential population, and in estimates of the star formation rate. However, the X-ray active population in the SMC during the same set of observations is very similar in number to that of the Magellanic Bridge. In the SMC, only four out of the

Table 2. Candidate sources in the Magellanic Bridge. Columns 5 and 6 indicate the revolutions that were mosaicked for each candidate detection. For IBIS detections the fluxes are given in the 15–35 keV range, while JEM-X fluxes are quoted in the 3–10 keV range.

Name	RA (J2000.0) (h m s)	Dec. (J2000.0) (d m s)	Error 90 per cent	IBIS revs	JEM-X revs	Flux (error) $\text{erg cm}^{-2} \text{s}^{-1}$
IGR J02048–7315	02 04 49	–73 15 27	1.8 arcmin		754–756	$1.7(0.7) \times 10^{-12}$
IGR J02220–7558	02 22 01	–75 57 59	1.8 arcmin		755–756	$3.5(1.1) \times 10^{-12}$
IGR J03144–7404	03 14 23	–74 04 23	4.9 arcmin	746–747		$4.4(0.9) \times 10^{-11}$




Figure 9. IBIS exposure map for the sum of all observations presented in this paper. The exposure on the eastern edge of the bridge is almost half of that on the western edge (SMC side). The image size is $10^\circ \times 15^\circ$, and the contours once again show the H I distribution throughout the SMC and Magellanic Bridge.

~60 HMXBs were X-ray active during these observations, so by observational comparison with the SMC, we may expect a population significantly larger than six Be/X-ray binaries in the Bridge.

6 CONCLUSIONS

In this paper, we presented two new hard X-ray sources in the Magellanic Bridge identified through wide-field, hard X-ray imaging. We described their timing and spectral characteristics and identified them as likely HMXBs located at the distance to the Magellanic Bridge. This puts the number of HMXBs in the Magellanic Bridge at three, with a further three candidate sources to add to the emerging population. Optical observations of this region show a large number of young stars, especially towards the western edge, consistent with a picture of in situ star formation as a result of tidal interaction between the SMC and LMC. The short-lived HMXBs presented here echo this scenario for star formation in the Bridge.

ACKNOWLEDGMENTS

Based on observations with *INTEGRAL*, an ESA project with instruments and science data centre funded by ESA members states (especially the PI countries: Denmark, France, Germany, Italy, Switzerland, Spain), Czech Republic and Poland, and with the participation of Russia and the USA. LJT wishes to thank the University of Southampton, whose support has made this research possible.

REFERENCES

- Bica E. L. D., Schmitt H. R., 1995, *ApJS*, 101, 41
 Coe M. J., 2005, *MNRAS*, 358, 1379
 Coe M. J., Edge W. R. T., Galache J. L., McBride V. A., 2005, *MNRAS*, 356, 502
 Coe M. J., McBride V. A., Bird A. J., Corbet R. H. D., 2008, *Astron. Tel.*, 1882, 1
 Coe M. J. et al., 2009, in van Loon J. T., Oliveira J. M., eds, *Proc. IAU Symp. Vol. 256, The Magellanic System: Stars, Gas, and Galaxies*. Cambridge Univ. Press, Cambridge, p. 367
 Demers S., Irwin M. J., 1991, *A&AS*, 91, 171
 Gardiner L. T., Hatzidimitriou D., 1992, *MNRAS*, 257, 195
 Gardiner L. T., Noguchi M., 1996, *MNRAS*, 278, 191
 Goldwurm A. et al., 2003, *A&A*, 411, L223
 Grimm H.-J., Gilfanov M., Sunyaev R., 2003, *MNRAS*, 339, 793
 Harris J., 2007, *ApJ*, 658, 345
 Harris J., Zaritsky D., 2004, *AJ*, 127, 1531
 Irwin M. J., Kunkel W. E., Demers S., 1985, *Nat*, 318, 160
 Kahabka P., Hilker M., 2005, *A&A*, 435, 9
 McGowan K. E. et al., 2007, *MNRAS*, 376, 759
 Mizuno N., Muller E., Maeda H., Kawamura A., Minamidani T., Onishi T., Mizuno A., Fukui Y., 2006, *ApJ*, 643, L107
 Monet D. G. et al., 2003, *AJ*, 125, 984
 Putman M. E., Staveley-Smith L., Freeman K. C., Gibson B. K., Barnes D. G., 2003, *ApJ*, 586, 170
 Ubertini P. et al., 2003, *A&A*, 411, L131
 Voges W. et al., 1999, *A&A*, 349, 389
 Voges W. et al., 2000, *IAU Circ.* 7432, 1
 Yokogawa J., Imanishi K., Tsujimoto M., Koyama K., Nishiuchi M., 2003, *PASJ*, 55, 161

This paper has been typeset from a \LaTeX file prepared by the author.